# MEDICATION THAT MUST BE STOPPED BEFORE SURGICAL PROCEDURE

***This list is may not include all medications – If you are unsure about a medication you take please contact your physician or pharmacy.***

## ALL INJECTABLE GLP-1 MEDICATIONS MUST BE STOPPED A FULL WEEK (7) DAYS BEFORE YOUR PROCEDURE

**GLP-1 Weight Loss & Diabetes (Dual/Multiple Use)**

- **Tirzepatide**
- **Zepbound**
- **Mounjaro**
- **Semaglutide**
- **Wegovy**
- **Liraglutide**
- **Saxenda**
- **Victoza**

**Type 2 Diabetes Only**

- **Dulaglutide** (Brand name: **Trulicity**)
- **Exenatide** (Brand names: **Byetta** twice-daily, **Bydureon BCise** weekly)
- **Lixisenatide** (Brand name: **Adlyxin**)

**Retatrutide** – A GLP-!, GIP, Glucagon "triple agonist"

****Oral GLP-1 Medication – Must be stopped one full day prior to your procedure AND you must be on clear liquids only for a full day prior to your procedure.**

- **Semaglutide (Oral)** -- Brand name: **Rybelsus** – *A daily tablet rather than an injection*
- **Semaglutide (Oral)** -- Brand name**: Wegovy** – *A daily tablet rather than an injection*
- **Orforglipron (Oral)** – Brand name: **Foundayo** – *A daily tablet rather than an injection*
-

## ALL SGLT-2 MEDICATIONS MUST BE STOPPED A FULL THREE (3) DAYS PRIOR TO YOUR PROCEDURE

### SGLT-2 MEDICATIONS

**Standalone Medications**

- **Canagliflozin** – Brand name: *Invokana*
- **Dapagliflozin** – Brand name: *Farxiga*
- **Empagliflozin** – Brand name: *Jardiance*
- **Ertugliflozin** – Brand name: *Steglatro*
- **Bexagliflozin** – Brand name: *Brenzavvy*

**Common Combination Medications**

- **Invokamet** (Canagliflozin + Metformin)
- **Xigduo XR** (Dapagliflozin + Metformin)
- **Synjardy** (Empagliflozin + Metformin)
- **Segluromet** (Ertugliflozin + Metformin)
- **Glyxambi** (Empagliflozin + Linagliptin)

## PHENTERMINE – MUST BE STOPPED A FULL FOUR (4) DAYS PRIOR TO YOUR PROCEDURE